

Estimation of Tropospheric Fluctuations using GPS Data

C.J. Naudet, (Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109; 818-354-2053 e-mail: cjn@logos.nasa.jpl.gov; Sponsor: O. J. Sovers)

Tropospheric turbulence is the limiting error source for Ka-band radio experiments and it limits coherence of high frequency radio interferometry. We discuss a method of extracting tropospheric fluctuation statistics from the GPS FLINN database. This database, an ongoing IGS effort at JPL, since 1992, produces daily solutions for orbits, receiver locations, clock offsets, tropospheric delays and residuals. GPS data is obtained in all meteorological conditions at over 30 sites with excellent temporal resolution.

It is assumed that the tropospheric fluctuations are well understood by the frozen flow Kolmogorov turbulence model and the hypothesis is formed that the GPS residuals are dominated by tropospheric delay fluctuations. The elevation and site dependence of the GPS residuals are discussed and are in agreement with the expectations of troposphere delay fluctuations. The GPS residuals are shown to have temporal statistics that are consistent with the assumed turbulence model. The turbulence parameter (Cn) is extracted from the GPS residuals and the absolute Cn magnitudes are found consistent with both WVR and VLB measurements. The seasonal and diurnal Cn dependence is presented and compared with a simple wet refractivity model.

1. 1995 Spring Meeting

2. 001300535

3. (a) C.J. Naudet
Ms 238-700
Jet Propulsion Lab.
4800 Oak Grove Dr.
Pasadena, CA 91109

(1) Tel: 818-354-2053

(c) Fax: 818-3934965

4. C

5. (a) None

(1) 0365 Troposphere
6964 Radio Wave propagation
3379 Turbulence

6. None

7. None

8. Charge \$60.00 to Visa Card
5437-0005-0673-7735 exp 12/96

9. C

10. None

11. No